

NEW ZEALAND DATA SHEET

1 Olimel® and PeriOlimel® (emulsions for infusion)

PeriOlimel N4-600E

Olimel N5-860E

Olimel N7-960, Olimel N7-960E

Olimel N9-840, Olimel N9-840E

Olimel N12-640 and Olimel N12-640E.

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Composition

Olimel/PeriOlimel with Electrolytes

Alanine, arginine, aspartic acid, glutamic acid, glycine, histidine, isoleucine, leucine, lysine acetate, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, sodium acetate trihydrate, sodium glycerophosphate hydrate, potassium chloride, magnesium chloride hexahydrate, calcium chloride dihydrate, glucose monohydrate, olive oil and soya oil.

Olimel without Electrolytes

Alanine, arginine, aspartic acid, glutamic acid, glycine, histidine, isoleucine, leucine, lysine acetate, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, glucose monohydrate, olive oil and soya oil.

Olimel/PeriOlimel is presented in the form of a 3-compartment bag. The individual compartments contain a glucose solution, a lipid emulsion and an amino acid solution (with/without electrolytes). In products containing electrolytes, calcium is included in the glucose solution compartment and other electrolytes are in the amino acid solution compartment. There are 7 different formulations of Olimel (4 with electrolytes and 3 without electrolytes) and one formulation of PeriOlimel. The general composition of the formulations are summarised below:

With electrolytes (E)	Without electrolytes	Nitrogen ¹	Amino acid solution ²	Glucose solution ³	Lipid emulsion ⁴
PeriOlimel N4-600E	-	4.0g/L	6.3%	18.75%	15.0%
Olimel N5-860E	-	5.2g/L	8.2%	28.75%	20.0%
Olimel N7-960E	Olimel N7-960	7.0g/L	11.1%	35.0%	20.0%
Olimel N9-840E	Olimel N9-840	9.0g/L	14.2%	27.5%	20.0%
Olimel N12-640E	Olimel N12-640	12.0g/L	14.2	27.5 %	17.5%

¹ prefixes N4, N5, N7, N9 and N12 refer to approx. nitrogen content in g/L.

² contains 17 amino acids (and electrolytes if present).

³ contains calcium if present.

⁴ contains refined olive oil (80%) and soya oil (20%).

Olimel/PeriOlimel contain egg lecithin. For the detailed formulations, refer to Appendix 1.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Emulsion for infusion (intravenous).

Appearance before reconstitution

- The amino acid and glucose solutions are clear and colourless or slightly yellow.
- The lipid emulsion is a homogeneous liquid with a milky appearance.

NEW ZEALAND DATA SHEET

Appearance after reconstitution:

After reconstitution/mixing of the contents of the 3 compartments, **Olimel/PeriOlimel** is a milk-like homogeneous liquid. The composition of the 3-in-1 admixture for each of the bag presentations are provided in Appendix 1.

Osmolarity

Olimel/PeriOlimel is a hypertonic emulsion. The osmolarity, osmolality and energy contents of the formulations are as follows:

Olimel/PeriOlimel	N4-600E	N5-860E	N7-960E	N7-960	N9-840E	N9-840	N12-640E	N12-640
Osmolarity approx. (mOsm/L)	760	1120	1360	1220	1310	1170	1270	1130
Energy content approx. (kcal/L)	700	990	1140	1140	1070	1070	950	950
Osmolality approx. (mOsm/kg water)	860	1340	1680	1490	1580	1410	1470	1310

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Olimel/PeriOlimel is indicated for parenteral nutrition for adults when oral or enteral nutrition is impossible, insufficient, or contraindicated.

4.2 Dose and method of administration

Use in one patient on one occasion only. It is recommended that after opening the bag, the contents should be used immediately, and should not be stored for subsequent infusion.

Due to its low osmolarity (760mOsmol/L), **PeriOlimel** N4-600E can be administered through a peripheral or central vein. Due to its high osmolarity (1120 – 1360mOsmol/L), **Olimel** N5-860E, N7-960E, N7-960, N9-840E, N9-840, N12-640E and N13-640 must only be administered through a central vein.

Although there is a natural content of trace elements and vitamins in the product, the levels are insufficient to meet body requirements. Trace elements and vitamins should be added in sufficient quantities to meet individual patient requirements and to prevent deficiencies from developing.

Adults

The dosage depends on the patient's energy expenditure, clinical status, body weight and ability to metabolise constituents of **Olimel/PeriOlimel**, as well as on additional energy or proteins given orally/enterally. Thus, the bag size should be then chosen with regard to the patient's body weight.

The average daily requirements for adults are:

- *Protein:* 0.16 to 0.35g nitrogen/kg body weight (1 to 2g of amino acids/kg) depending on the patient's nutritional status and degree of catabolic stress. Special populations may require up to 0.4g nitrogen/kg body weight (2.5g of amino acids/kg)
- *Energy:* 20 to 40kcal/kg.
- *Fluid:* 20 to 40mL fluid/kg, or 1 to 1.5mL per expended kcal.

Maximum daily dose

The maximum daily dose varies with the clinical condition of the patient and may even change from day to day.

PeriOlimel N4-600E

For **PeriOlimel**, the maximum daily dose is defined by fluid intake, 40mL/kg, corresponding to 1g/kg amino acids, 3g/kg glucose, 1.2g/kg lipids, 0.8mmol/kg sodium, and 0.6mmol/kg potassium.

NEW ZEALAND DATA SHEET

For example, for a 70kg patient, this would be equivalent to 2,800mL **PeriOlimel** per day, resulting in an intake of 71g amino acids, 210g glucose, and 84g lipids (i.e., 1,680 non-protein kcal and 1,960 total kcal).

Olimel N5-860E

For **Olimel** N5E, the maximum daily dose is defined by fluid intake, 40mL/kg, corresponding to 1.3g/kg amino acids, 4.6g/kg glucose, 1.6g/kg lipids, 1.4mmol/kg sodium, and 1.2mmol/kg potassium.

For example, for a 70kg patient, this would be equivalent to 2,800mL **Olimel** per day, resulting in an intake of 92g amino acids, 322g glucose, and 112g lipids (i.e., 2,408 non-protein kcal and 2,772 total kcal).

Olimel N7-960

For **Olimel** N7, the maximum daily dose is defined by total caloric intake, 40kcal/kg provided in a volume of 35mL/kg, corresponding to 1.5g/kg amino acids, 4.9g/kg glucose, and 1.4g/kg lipids.

For example, for a 70kg patient, this would be equivalent to 2,450mL **Olimel** per day, resulting in an intake of 108g amino acids, 343g glucose, and 98g lipids (i.e., 2,352 non-protein kcal and 2,793 total kcal).

Olimel N7-960E

For **Olimel** N7E, the maximum daily dose is defined by total caloric intake, 40kcal/kg provided in a volume of 35mL/kg, corresponding to 1.5g/kg amino acids, 4.9g/kg glucose, 1.4g/kg lipids, 1.2mmol/kg sodium, and 1.1mmol/kg potassium.

For example, for a 70kg patient, this would be equivalent to 2,450mL **Olimel** per day, resulting in an intake of 108g amino acids, 343g glucose, and 98g lipids (i.e., 2,352 non-protein kcal and 2,793 total kcal).

Olimel N9-840

For **Olimel** N9, the maximum daily dose defined by amino acids intake, 35mL/kg corresponding to 2.0g/kg amino acids, 3.9g/kg glucose, and 1.4g/kg lipids.

For example, for a 70kg patient, this would be equivalent to 2,450mL **Olimel** per day, resulting in an intake of 140g amino acids, 270g glucose, and 98g lipids (i.e., 2,058 non-protein kcal and 2,622 total kcal).

Olimel N9-840E

For **Olimel** N9E, the maximum daily dose is defined by amino acids intake, 35mL/kg corresponding to 2.0g/kg amino acids, 3.9g/kg glucose, 1.4g/kg lipids, 1.2mmol/kg sodium, and 1.1mmol/kg potassium.

For example, for a 70kg patient, this would be equivalent to 2,450mL **Olimel** per day, resulting in an intake of 140g amino acids, 270g glucose, and 98g lipids (i.e., 2,058 non-protein kcal and 2,622 total kcal).

Olimel N12-640

For **Olimel** N12-640, the maximum daily dose defined by amino acids intake, 26mL/kg corresponding to 2.0g/kg amino acids, 1.9g/kg glucose, and 0.9g/kg lipids.

For example, for a 70kg patient, this would be equivalent to 1,820mL **Olimel** per day, resulting in an intake of 138g amino acids, 133g glucose, and 64g lipids (i.e., 1,171 non-protein kcal and 1,723 total kcal).

Olimel N12-640E

For **Olimel** N12-640E, the maximum daily dose defined by amino acids intake, 26mL/kg corresponding to 2.0g/kg amino acids, 1.9g/kg glucose, 0.9g/kg lipids, 0.9mmol/kg sodium, and 0.8mmol/kg potassium.

For example, for a 70kg patient, this would be equivalent to 1,820mL **Olimel** per day, resulting in an intake of 138g amino acids, 133g glucose, and 64g lipids (i.e., 1,171 non-protein kcal and 1,723 total kcal).

NEW ZEALAND DATA SHEET

In Continuous Renal Replacement Therapy (CRRT) and patients with morbid obesity

For **Olimesel** N12-640, the maximum daily dose is defined by amino acids intake, 33mL/kg corresponding to 2.5g/kg amino acids, 2.4g/kg glucose, 1.2g/kg lipids.

For **Olimesel** N12-640E, the maximum daily dose is defined by amino acids intake, 33mL/kg corresponding to 2.5g/kg amino acids, 2.4g/kg glucose, 1.2g/kg lipids, 1.2mmol/kg sodium, and 1.0mmol/kg potassium.

In patients with morbid obesity, the dosage should be calculated on basis of the Ideal Body Weight (IBW).

For example, for a 70kg patient, this would be equivalent to 2,310mL **Olimesel** N12-640E/N12-640 per day, resulting in an intake of 175g amino acids, 169g glucose, and 81g lipids (i.e., 1,486 non-protein kcal and 2,187 total kcal).

The maximum daily dose should not be exceeded. Due to the static composition of the multi-chamber bag, the ability to simultaneously meet all nutrient needs of the patient may not be possible. Clinical situations may exist where patients require amounts of nutrients varying from the composition of the static bag. In this situation the impact of any volume (dose) adjustments must be taken into consideration and the resultant effect this will have on the dosing of all other nutrient components of **Olimesel**.

The flow rate should be increased gradually during the first hour. The administration flow rate must be adjusted taking into account the dose being administered, the daily volume intake, and the duration of the infusion.

PeriOlimesel N4-600E

For **PeriOlimesel** N4E, the maximum infusion rate is 3.2mL/kg/hour, corresponding to 0.08g/kg/hour amino acids, 0.24g/kg/hour glucose, and 0.10g/kg/hour lipids.

Olimesel N5-860E

For **Olimesel** N5E, the maximum infusion rate is 2.1mL/kg/hour, corresponding to 0.07g/kg/hour amino acids, 0.24g/kg/hour glucose, and 0.08g/kg/hour lipids.

Olimesel N7-960E/N7-960

For **Olimesel** N7E/N7, the maximum infusion rate is 1.7 mL/kg/hour, corresponding to 0.08 g/kg/hour amino acids, 0.24 g/kg/hour glucose, and 0.07 g/kg/hour lipids.

Olimesel N9-840E/N9-840

For **Olimesel** N9E/N9, the maximum infusion rate is 1.8mL/kg/hour, corresponding to 0.10g/kg/hour amino acids, 0.19g/kg/hour glucose, and 0.07g/kg/hour lipids.

Olimesel N12-640E/N12-640

For **Olimesel** N12-640/N12-640E, the maximum infusion rate is 1.3mL/kg/hour, corresponding to 0.10g/kg/hour amino acids, 0.10g/kg/hour glucose, and 0.05g/kg/hour lipids.

The recommended duration of infusion for a parenteral nutrition bag is between 12 and 24 hours. Treatment with parenteral nutrition may be continued for as long as is required by the patient's condition.

Method of preparation

Before opening the overpouch, check the colour of the oxygen indicator. Compare it to the reference colour printed next to the OK symbol and depicted in the printed area of the indicator label. Do not use the product if the colour of the oxygen indicator does not correspond to the reference colour printed next to OK symbol.

Preparation for administration

a) To open

Remove the protective overpouch.

Discard the oxygen absorber / oxygen indicator sachet.

Confirm the integrity of the bag and of the non-permanent seals.

NEW ZEALAND DATA SHEET

Use only if the bag is not damaged, if the non-permanent seals are intact (i.e. no mixture of the contents of the three compartments), if the amino acids solution and the glucose solution are clear, colourless or slightly yellow, practically free of visible particles, and if the lipid emulsion is a homogeneous liquid with a milky appearance.

b) Mixing the solutions and the emulsion

Ensure that the product is at room temperature when breaking the non-permanent seals.

Manually roll the bag onto itself, starting at the top of the bag (hanger end). The non-permanent seals will disappear from the side near the inlets. Continue to roll until the seals are open along approximately half of their length.

Mix by inverting the bag at least 3 times.

After reconstitution, the mixture is a homogeneous emulsion with a milky appearance.

c) Additions

The capacity of the bag is sufficient to enable additions such as vitamins, electrolytes and trace elements.

Any addition (including vitamins) may be made into the reconstituted mixture (after the non-permanent seals have been opened and after the contents of the three compartments have been mixed). Vitamins may also be added into the glucose compartment before the mixture is reconstituted (before opening the non-permanent seals and before mixing the 3 compartments).

When making additions to the formulation, the final osmolarity of the mixture should be measured before administration via a peripheral vein.

The maximum total levels of sodium, magnesium, potassium and calcium listed in the table below were demonstrated by stability data and should not be considered dosage recommendations.

Electrolyte supplementation should be dictated by the patients clinical needs and should not exceed nutritional guidelines. When making additions to formulations containing electrolytes, the amount of electrolytes already present in the bag should be taken into account.

Additions must be performed by qualified personnel under aseptic conditions.

Olimel/PeriOlimel may only be added to medicinal or nutritional solutions for which compatibility has been documented.

PeriOlimel N4-600E may be supplemented with electrolytes according to the following table:

Additions to PeriOlimel N4-600E per 1000mL			
	Included level	Maximal further addition	Maximal total level
Sodium	21.0mmol	129.0mmol	150.0mmol
Potassium	16.0mmol	134.0mmol	150.0mmol
Magnesium	2.2mmol	3.4mmol	5.6mmol
Calcium	2.0mmol	3.0mmol	5.0mmol
Phosphate	8.5mmol(*)	Inorganic Phosphate 8.0mmol	16.5mmol(*)
		Organic Phosphate 15.0mmol	23.5mmol(*)
(*) including phosphate provided by the lipid emulsion			

The Olimel formulations containing electrolytes

Olimel N5-860E, N7-960E and N9-840E may be supplemented with electrolytes according to the table below:

NEW ZEALAND DATA SHEET

Additions to Olimel N5-860E, N7-960E and N9-840E per 1000mL			
	Included level	Maximal further addition	Maximal total level
Sodium	35.0mmol	115.0mmol	150.0mmol
Potassium	30.0mmol	120.0mmol	150.0mmol
Magnesium	4.0mmol	1.6mmol	5.6mmol
Calcium	3.5mmol	1.5mmol	5.0mmol
Phosphate	15mmol(#)	Inorganic Phosphate 3.0mmol	18.0mmol(#)
		Organic Phosphate 10.0mmol	25.0mmol(#)
(#) including phosphate provided by the lipid emulsion			

The *Olimel* formulations without electrolytes

Olimel N7-960 and N9-840 may be supplemented with electrolytes according to the following table:

Additions to Olimel N7-960 and N9-840 per 1000mL			
	Included level	Maximal further addition	Maximal total level
Sodium	0mmol	150.0mmol	150.0mmol
Potassium	0mmol	150.0mmol	150.0mmol
Magnesium	0mmol	5.6mmol	5.6mmol
Calcium	0mmol	5.0mmol	5.0mmol
Phosphate	3mmol(+)	Inorganic Phosphate 8.0mmol	11.0mmol(+)
		Organic Phosphate 22.0mmol	25.0mmol(+)
(+) including phosphate provided by the lipid emulsion			

The *Olimel* formulations - N12-640 and N12-640E

Olimel N12-640 and N12-640E may be supplemented with electrolytes, inorganic/organic phosphate according to the following tables:

Possible supplementations for 1000 mL Olimel N12E			
	Included level	Maximal further addition	Maximal total level
Sodium	35mmol	115mmol	150mmol
Potassium	30mmol	120mmol	150mmol
Magnesium	4.0mmol	1.6mmol	5.6mmol
Calcium	3.5mmol	1.5mmol	5.0mmol
Inorganic Phosphate	0mmol	10mmol Pi or	10 mmolPi + 15mmol Po or
Organic Phosphate	15mmol a	10 mmol Po ^b	25mmol Po ^{a,b}
^a Including Phosphate provided by the lipid emulsion ^b Pi - inorganic phosphate; Po - organic phosphate			

NEW ZEALAND DATA SHEET

Possible supplementations for 1000 mL Olimel N12			
	Included level	Maximal further addition	Maximal total level
Sodium	0mmol	150mmol	150mmol
Potassium	0mmol	150mmol	150mmol
Magnesium	0mmol	5.6mmol	5.6mmol
Calcium	0mmol	5.0mmol	5.0mmol
Inorganic Phosphate	0mmol	10mmol Pi + 12mmol Po or 22mmol Po ^b	10mmol Pi + 15mmol Po or 25mmol Po ^{a,b}
Organic Phosphate	3mmol ^a		
^a Including Phosphate provided by the lipid emulsion ^b Pi - inorganic phosphate; Po - organic phosphate			

Trace elements and vitamins

Stability has been demonstrated with commercially available preparations of vitamins and trace elements (containing up to 1mg of iron).

For further compatibility information with the different products and storage conditions of the different admixtures please contact Baxter Healthcare.

To perform an addition:

- Aseptic conditions must be observed.
- Prepare the injection site of the bag.
- Puncture the injection site and inject the additives using an injection needle or a reconstitution device.
- Mix content of the bag and the additives.

d) Preparation of the infusion

Aseptic conditions must be observed.

Suspend the bag.

Remove the plastic protector from the administration outlet.

Firmly insert the spike of the infusion set into the administration outlet.

e) Administration

Use in one patient on one occasion only.

Only administer the product after the non-permanent seals between the three compartments have been broken and the contents of the three compartments have been mixed. Ensure that the final emulsion for infusion does not show any evidence of phase separation.

After opening the bag, the content must be used immediately, should not be stored for a subsequent infusion. Do not reconnect any partially used bag.

Do not connect in series in order to avoid the possibility of air embolism due to gas contained in the first bag. Any unused product or waste material and all necessary disposable devices must be discarded.

4.3 Contraindications

Use of **Olimel/PeriOlimel** is contraindicated in the following situations:

- In premature neonates, infants and children less than 2 years old.
- Known hypersensitivity to egg or soya proteins, peanut protein, corn (maize) and corn products, components of the container, or to any of the ingredients including active substances and/or excipients.
- Congenital abnormalities of amino acid metabolism.
- Severe hyperlipidaemia or severe disorders of lipid metabolism characterised by hypertriglyceridaemia.
- Severe hyperglycaemia.

NEW ZEALAND DATA SHEET

- Unstable conditions (for example, following severe post-traumatic conditions, acute phase of circulatory shock, acute myocardial infarction, severe sepsis and hyperosmolar coma).
- **Olimel/PeriOlimel** formulations with electrolytes must not be administered to patients with pathologically elevated plasma concentrations of sodium, potassium, magnesium, calcium and/or phosphorus.

Use with caution in patients with severe liver insufficiency, including cholestasis or elevated liver enzymes. Liver function parameters should be closely monitored.

4.4 Special warnings and precautions for use

Allergic reactions

The infusion must be stopped immediately if any signs or symptoms of an allergic reaction (such as fever, shivering, skin rashes or dyspnoea) develop.

Olimel/PeriOlimel contain glucose. Solutions containing glucose should be used with caution in patients with known allergy to corn or corn products.

Pulmonary

Pulmonary vascular precipitates causing pulmonary vascular emboli and pulmonary distress have been reported in patients receiving parenteral nutrition. In some cases, fatal outcomes have occurred. Excessive addition of calcium and phosphate increases the risk of the formation of calcium phosphate precipitates. Precipitates have been reported even in the absence of phosphate salt in the solution. Suspected precipitate formation in the blood stream have also been reported.

In addition to inspection of the solution, the infusion set and catheter should also periodically be checked for precipitates.

If signs of pulmonary distress occur, the infusion should be stopped and medical evaluation initiated.

Compatibility

No additions to the bag should be made without first checking the compatibility, as formation of precipitates or destabilisation of the lipid emulsion could result in vascular occlusion (see section 4.5).

Ceftriaxone must not be administered simultaneously with intravenous calcium-containing solutions, including **Olimel/PeriOlimel**, through the same infusion line (e.g., via Y-connector) because of the risk of precipitation of ceftriaxone-calcium salt.

If the same infusion line is used for sequential administration, the line must be thoroughly flushed with a compatible fluid between infusions.

Infection and sepsis

Infection and sepsis may occur as a result of the use of intravenous catheters to administer parenteral formulations, poor maintenance of catheters or contaminated solutions. Immunosuppression by medicines and other factors such as hyperglycaemia, malnutrition and/or their underlying disease state may predispose patients to infectious complications. Careful symptomatic and laboratory monitoring for fever/chills, leukocytosis, technical complications with the access device, and hyperglycaemia can help recognise early infections. The occurrence of septic complications can be decreased with heightened emphasis on aseptic technique in catheter placement, maintenance, as well as aseptic technique in nutritional formula preparation.

Fat overload syndrome

“Fat overload syndrome” has been reported with similar products. This may be caused by inappropriate administration (e.g. overdose and/or infusion rate higher than recommended, see section 4.9); however, the signs and symptoms of this syndrome may also occur when the product is administered according to instructions. The reduced or limited ability to metabolise the lipids contained in **Olimel/PeriOlimel** accompanied by prolonged

NEW ZEALAND DATA SHEET

plasma clearance may result in a fat overload syndrome. This syndrome is associated with a sudden deterioration in the patient's clinical condition and is characterised by findings such as fever, anaemia, leucopenia, thrombocytopenia, coagulation disorders, hyperlipidaemia, liver fatty infiltration (hepatomegaly), deteriorating liver function, and central nervous system manifestations (e.g. coma). The syndrome is usually reversible when the infusion of the lipid emulsion is stopped.

Refeeding syndrome

Refeeding severely undernourished patients may result in the refeeding syndrome that is characterised by the shift of potassium, phosphorus and magnesium intracellularly as the patient becomes anabolic. Thiamine deficiency and fluid retention may also develop. Careful monitoring and slowly increasing nutrient intakes while avoiding overfeeding can prevent these complications. This syndrome has been reported with similar products.

Preparation and administration

If the final mixture is hypertonic, it may cause irritation of the vein when administered into a peripheral vein.

While **PeriOlimel** N4-600E may be administered through a peripheral vein, thrombophlebitis may develop. The catheter insertion site must be monitored daily for local signs of thrombophlebitis.

Olimel N5-860E, N7-960, N7-960E, N9-840, N9-840E, N12-640 and N12-640E must only be administered through a central vein.

Do not connect bags in series in order to avoid air embolism due to possible residual air contained in the primary bag.

Do not add other medicinal products or substances to one of the three compartments of the bag or to the reconstituted solution/emulsion without firstly confirming their compatibility and the stability of the resulting preparation (in particular stability of the lipid emulsion).

Extravasation

Extravasation has been reported with the administration of **Olimel/PeriOlimel**. Catheter site should be monitored regularly to identify signs of extravasation.

If extravasation occurs the administration should be stopped immediately, keeping the inserted catheter or cannula in place for immediate management of the patient. If possible, aspiration should be performed through the inserted catheter/cannula in order to reduce the amount of fluid present in the tissues before removing the catheter/cannula.

Depending on the extravasated product (including the product(s) being mixed with **Olimel**, if applicable) and the stage/extent of any injury, appropriate specific measures should be taken. Options for management may include non-pharmacologic, pharmacologic and/or surgical intervention. In case of large extravasation, plastic surgeon advice should be sought within the first 72 hours.

The extravasation site should be monitored at least every 4 hours during the first 24 hours, then once daily.

The infusion should not be restarted in the same central vein.

Monitoring

Monitor water and electrolyte balance, serum osmolarity, serum triglycerides, acid/base balance, blood glucose, liver and kidney function, and blood count, including platelets and coagulation parameters throughout treatment.

Hypercalciuria may occur in high-protein dose PN, and patients should be monitored for metabolic consequences. In particular, calcium and phosphate levels should be monitored.

NEW ZEALAND DATA SHEET

In addition, regular clinical and laboratory tests are required particularly in cases of:

- Amino acid metabolism disorders (see section 4.3).
- Hepatic insufficiency because of the risk of developing or worsening neurological disorders associated with hyperammonaemia.
- Renal insufficiency, particularly if hyperkalaemia is present; risk of developing or worsening metabolic acidosis and hyperazotaemia if extra-renal waste removal is not being performed.
- Metabolic acidosis (administration of carbohydrates is not recommended in the presence of lactic acidosis).
- Diabetes mellitus: monitoring of glucose concentrations, glucosuria, ketonuria and, where applicable, adjustment of insulin dosages.
- Coagulation disorders.
- Anaemia.
- Hyperlipidaemia (because of the presence of lipids in the emulsion for infusion).

The blood count and coagulation factors must be monitored more carefully during long-term administration (several weeks).

Cardiovascular

Use with caution in patients with pulmonary oedema or heart failure. Fluid status should be closely monitored.

Endocrine and metabolism

Metabolic complications may occur if the nutrient intake is not adapted to the patient's requirements, or the metabolic capacity of any given dietary component is not accurately assessed. Adverse metabolic effects may arise from administration of inadequate or excessive nutrients or from inappropriate composition of an admixture for a particular patient's needs.

Serum triglyceride concentrations and the ability of the body to metabolise lipids must be checked regularly. If a lipid metabolism abnormality is suspected, monitoring of serum triglycerides is recommended as clinically necessary.

In the event of hyperglycaemia, the infusion rate of **Olimel/PeriOlimel** must be adjusted and/or insulin administered.

Hepatobiliary disorders

Hepatobiliary disorders including cholestasis, hepatic steatosis, fibrosis and cirrhosis, possibly leading to hepatic failure, as well as cholecystitis and cholelithiasis are known to develop in some patients on parenteral nutrition. The etiology of these disorders is thought to be multifactorial and may differ between patients. Patients developing abnormal liver function parameters or other signs of hepatobiliary disorders should be assessed early by a clinician knowledgeable in liver diseases in order to identify possible causative and contributory factors, and possible therapeutic and prophylactic interventions.

Use in hepatic impairment

Use with caution in patients with hepatic insufficiency because of the risk of developing or worsening neurological disorders associated with hyperammonaemia. Regular clinical and laboratory tests are required, particularly liver function parameters, blood glucose, electrolytes and triglycerides.

Use in renal impairment

Use with caution in patients with renal insufficiency, particularly if hyperkalaemia is present, because of the risk of developing or worsening metabolic acidosis and hyperazotaemia if extra-renal waste removal is not being performed. Fluid, triglycerides and electrolyte status should be closely monitored in these patients.

Severe water and electrolyte equilibration disorders, severe fluid overload states, and severe metabolic disorders should be corrected before starting the infusion.

NEW ZEALAND DATA SHEET

Use in the elderly

In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or medication therapy. Due to the risk of azotaemia, appropriate monitoring (e.g. of urine urea nitrogen and blood urea nitrogen) should be considered in elderly patients.

Paediatric use

This product is contraindicated in premature neonates, infants and children less than 2 years old (see 4.3). There have been no studies performed in the paediatric population.

Effects on laboratory tests

The lipids contained in this emulsion may interfere with the results of certain laboratory tests (for example, bilirubin, lactate dehydrogenase, oxygen saturation, blood haemoglobin) if the blood sample is taken before the lipids are eliminated (these are generally eliminated after a period of 5 to 6 hours without receiving lipids).

4.5 Interaction with other medicines and other forms of interaction

No interaction studies have been performed with **Olimel/PeriOlimel**.

Do not add other medicinal products or substances to one of the three compartments of the bag or to the reconstituted solution/emulsion without firstly confirming their compatibility and the stability of the resulting preparation (in particular stability of the lipid emulsion or formation of precipitates).

As with any parenteral nutrition admixture, calcium and phosphate ratios must be considered. Excess addition of calcium and phosphate, especially in the form of mineral salts, may result in the formation of calcium phosphate precipitates.

Olimel/PeriOlimel must not be administered simultaneously with blood through the same infusion tubing because of the risk of pseudoagglutination.

Due to the risk of precipitation, **Olimel/PeriOlimel** should not be administered through the same infusion line or admixed together with ampicillin or fosphenytoin.

Olimel/PeriOlimel contains calcium ions which pose additional risk of coagulation precipitated in citrate anticoagulated/preserved blood or components. This only applies to products containing electrolytes.

Soya oil has a natural content of vitamin K1 that may counteract the anticoagulant activity of coumarin derivatives, including warfarin.

Due to the potassium content of **Olimel/PeriOlimel** (with electrolyte formulations), special care should be taken in patients simultaneously treated with potassium sparing diuretics with ACE inhibitors, angiotensin II receptor antagonists, or the immunosuppressants tacrolimus and cyclosporin in view of the risk of hyperkalaemia.

4.6 Fertility, pregnancy and lactation

Effect on fertility

No studies have been conducted to assess the effects of **Olimel/PeriOlimel** on fertility.

Pregnancy (Category - exempt)

There are no adequate data on use of **Olimel/PeriOlimel** in pregnant women. Physicians should carefully consider the potential risks and benefits for each specific patient before prescribing **Olimel/PeriOlimel**.

Breast-feeding

There are no adequate data on use of **Olimel/PeriOlimel** in lactating women. Following intravenous infusion, most of the active ingredients contained in **Olimel/PeriOlimel** are expected to be excreted in human milk and the safety

NEW ZEALAND DATA SHEET

of the breastfeeding infant has not been established. Physicians should carefully consider the potential risks and benefits for each specific patient before prescribing **Olimel/PeriOlimel**.

4.7 Effects on ability to drive and use machines

The effects of **Olimel/PeriOlimel** on a person's ability to drive and use machines were not assessed as part of its registration.

4.8 Undesirable effects

The safety and clinical efficacy of **Olimel** N9-840 was assessed in one double-blind randomised study with an active control over five days. Twenty-eight patients with different medical conditions (post-surgery fasting, severe malnutrition, enteral intake insufficient or forbidden) were included in the **Olimel** group and received the medicine at up to 40mL/kg/day.

The investigator judged the following seven adverse reactions as related to **Olimel**:

Clinical Trial Adverse Reactions	
System Organ Class (SOC)	Preferred MedDRA Term
Cardiac disorders	Tachycardia
Gastrointestinal disorders	Abdominal pain Diarrhoea Nausea
Metabolism and nutritional disorders	Decreased appetite Hypertriglyceridaemia
Vascular disorders	Hypertension

Post-marketing experience

The following adverse reactions have been reported in the Post-marketing experience, listed by MedDRA System Organ Class (SOC), then by Preferred Term in order of severity.

GASTROINTESTINAL DISORDERS: Vomiting

SKIN AND SUBCUTANEOUS SKIN DISORDERS: Rash

GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS: Injection site extravasation, Pyrexia, Chills.

The following adverse reactions have been reported with other similar products:

- Fat overload syndrome.
- Cholestasis, elevated liver enzymes and Azotaemia.
- Pulmonary vascular precipitates (pulmonary vascular emboli and pulmonary distress).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continuing monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://nzphv.otago.ac.nz/reporting/>

4.9 Overdose

In the event of inappropriate administration (overdose and/or infusion rate higher than recommended), nausea, vomiting, chills, headache, hot flush, hyperhidrosis and electrolyte disturbances and signs of hypervolaemia or acidosis may occur and result in severe or fatal consequences. In such situations the infusion must be stopped immediately. If medically appropriate, further intervention may be indicated.

Hyperglycaemia, glucosuria, and hyperosmolar syndrome may develop if glucose infusion rate exceeds clearance.

NEW ZEALAND DATA SHEET

In some serious cases, haemodialysis, haemofiltration, or haemodiafiltration may be necessary.

The reduced or limited ability to metabolise lipids may result in fat overload syndrome, the results of which are usually reversible after infusion of the lipid emulsion is stopped.

For advice on the management of overdose please contact the National Poisons Centre on phone number: 0800 764 766 [0800 POISON] in New Zealand (or 131126 in Australia).

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group

Solutions for parenteral nutrition/mixtures.

ATC code

B05 BA 10.

Mechanism of Action/Pharmacological actions

This is a 3-in-1 admixture enabling the nitrogen/energy balance to be maintained from the nitrogen source (L series amino acids) and energy in the form of glucose and essential fatty acids. Nitrogen and energy are required for normal functioning of all cells in the body, and are important for protein synthesis, growth, wound healing, immune function, muscle function, any other cellular activities.

The amino acids solution contains 17 amino acids (including 8 essential amino acids), which are required for protein synthesis. Amino acids also represent an energy source, their oxidation resulting in excretion of nitrogen in the form of urea. The amino acids profile is as follows:

- Essential amino acids/total amino acids: 44.8%.
- Branched-chain amino acids/total amino acids: 18.3%.

The formulations without electrolytes allow individual electrolyte intake to be adapted to meet specific requirements.

The lipid emulsion included in **Olimel/PeriOlimel**, is an association of refined olive oil and refined soya oil (ratio 80/20), with the following approximate distribution of fatty acids:

- 15% saturated fatty acids (SFA).
- 65% monounsaturated fatty acids (MUFA).
- 20% polyunsaturated essential fatty acids (PUFA).

The phospholipid/triglyceride ratio is 0.06. The moderate essential fatty acid (EFA) content improves the status of their upper derivatives while correcting EFA deficiency.

Olive oil contains significant amounts of alpha-tocopherol, when combined with a moderate PUFA intake, contributes to improve vitamin E status and reduce lipid peroxidation.

The carbohydrate source is glucose. Glucose is the primary source of energy in the body.

Molecular formulae and CAS numbers

The molecular formula and CAS (Chemical Abstract Service) registry number of the active substances are listed in the following table.

NEW ZEALAND DATA SHEET

<i>Molecular formula and CAS registry number of the active substances</i>		
Active substances	Molecular Formula	CAS Number
L-Alanine	C ₃ H ₇ NO ₂	56-41-7
L-Arginine	C ₆ H ₁₄ N ₄ O ₂	74-79-3
L-Aspartic acid	C ₄ H ₇ NO ₄	56-84-8
L-Glutamic acid	C ₅ H ₉ NO ₄	56-86-0
Glycine	C ₂ H ₅ NO ₂	56-40-6
L-Histidine	C ₆ H ₉ N ₃ O ₂	71-00-1
L-Isoleucine	C ₆ H ₁₃ NO ₂	73-32-5
L-Leucine	C ₆ H ₁₃ NO ₂	61-90-5
L-Lysine acetate	C ₆ H ₁₄ N ₂ O ₂ ·C ₂ H ₄ O ₂	57282-49-2
L-Methionine	C ₅ H ₁₁ NO ₂ S	63-68-3
L-Phenylalanine	C ₉ H ₁₁ NO ₂	63-91-2
L-Proline	C ₅ H ₉ NO ₂	147-85-3
L-Serine	C ₃ H ₇ NO ₃	56-45-1
L-Threonine	C ₄ H ₉ NO ₃	72-19-5
L-Tryptophan	C ₁₁ H ₁₂ N ₂ O ₂	73-22-3
L-Tyrosine	C ₉ H ₁₁ NO ₃	60-18-4
L-Valine	C ₅ H ₁₁ NO ₂	72-18-4
Sodium acetate, trihydrate	C ₂ H ₃ NaO ₂ ·3H ₂ O	6131-90-4
Sodium glycerophosphate hydrate	C ₃ H ₇ Na ₂ O ₆ P·xH ₂ O (degree of hydration: x = 4 to 6)	1334-74-3 (anhydrous)
Potassium chloride	KCl	7447-40-7
Magnesium chloride, hexahydrate	MgCl ₂ ·6H ₂ O	7791-18-6
Calcium chloride, dihydrate	CaCl ₂ ·2H ₂ O	10035-04-8
Glucose monohydrate	C ₆ H ₁₂ O ₆ ·H ₂ O	5996-10-1
Refined olive oil	Complex mixture of triglycerides; predominant fatty acids in olive oil are oleic, palmitic and linoleic.	8001-25-0
Refined soya oil	Complex mixture of triglycerides; predominant fatty acids in soya oil are linoleic, palmitic and linolenic.	8001-22-7

Clinical efficacy and safety

Study ICS1063B/P01/03/Mu.F was a prospective randomised double-blind multicenter study performed in fifty six hospitalized patients (age range 18 – 85 years) to evaluate safety and nutritional efficacy of **Olimel** N9-840 compared to **OliClinomel** N8-800 (not registered in New Zealand [or Australia] but contains the same ingredients as the **OliClinomel** products registered in NZ). The study was conducted in a variety of patients (primarily post-surgery and trauma) who required balanced parenteral nutrition representing at least 50% of the daily non-protein energy requirements for 5 days. The primary nutritional efficacy endpoint was transthyretin (pre-albumin) levels. Safety was evaluated using adverse events, vital signs, and biochemical markers for renal (urea, creatinine), hepatic (AST, ALT, alkaline phosphatase, GGT, bilirubin), haematologic (RBC count, haemoglobin, haematocrit, platelets, WBCs, lymphocytes, neutrophils, monocytes, eosinophils, basophils), organ functions as well as glucose and lipid parameters (triglycerides, cholesterol).

Efficacy analysis on the *per protocol* (PP) and *intent-to-treat* (ITT) populations showed no difference between the **Olimel** and **OliClinomel** groups on the primary endpoint (transthyretin), which improved from baseline to Day 5/end of treatment.

NEW ZEALAND DATA SHEET

Changes in Mean Pre-albumin (Transthyretin) Levels – Study ICS1063B/P01/03/Mu.F		
Treatment Group Study Population	Mean ± SD Transthyretin Levels (g/L)	
	Baseline	Day 5/EOT
Olimel N9-840		
Intent-to-Treat Population (n = 24)	0.144 ± 0.075	0.206 ± 0.142
Per Protocol Population (n = 24)	0.144 ± 0.075	0.206 ± 0.142
OliClinomel N8-800		
Intent-to-Treat Population (n = 26)	0.146 ± 0.083	0.181 ± 0.082
Per Protocol Population (n = 23)	0.139 ± 0.078	0.172 ± 0.080
EOT: End of treatment. SD: Standard deviation.		

The safety of the two formulations was comparable. There was no difference between the treatment groups for any clinical laboratory or vital sign parameters evaluated during the study.

5.2 Pharmacokinetic properties

The ingredients of the emulsion for infusion (amino acids, glucose and lipids) are distributed, metabolised and eliminated in the same way as if they had been administered individually.

The pharmacokinetic properties of the amino acids administered intravenously are principally the same as those of amino acids supplied by oral feeding. Amino acids from food proteins, however, first pass through the portal vein before reaching the systemic circulation.

The elimination rate of lipid emulsions depends on particle size. Small lipid particles appear to delay clearance whereas they increase lipolysis by lipoprotein lipase. Most of the lipid particle sizes are in the range of chylomicrons (0.08 – 0.6 micrometers) with the mean diameter of less than 0.35 micrometers. However, it may contain a small fraction (up to 2.5%) of particles having a diameter of more than 0.75 micrometer.

5.3 Preclinical safety data

Genotoxicity

No genotoxicity studies have been conducted with **Olimel/PeriOlimel**.

Carcinogenicity

No carcinogenicity studies have been conducted with **Olimel/PeriOlimel**.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Olimel/PeriOlimel contains the following excipients:

- Egg lecithin (purified egg phosphatide).
- Glycerol.
- Sodium oleate.
- Sodium hydroxide/Glacial acetic acid/Hydrochloric acid (for pH adjustment).
- Water for injections.

6.2 Incompatibilities

See section 4.5.

6.3 Shelf life

24 months from date of manufacture (stored at or below 25°C).

The expiry date can be found on the packaging.

NEW ZEALAND DATA SHEET

After reconstitution

7 days reconstituted stored at 2° to 8°C (Refrigerate, do not freeze), followed by 48 hours reconstituted stored at or below 25°C.

6.4 Special precautions for storage

Store below 25°C. Do not freeze. Store in overpouch.

After reconstitution

It is recommended that the product is used immediately after the non-permanent seals between the three compartments have been opened. However, the stability of the reconstituted emulsion has been demonstrated for 7 days (between 2°C and 8°C) followed by 48 hours at temperature not exceeding 25°C.

After addition of supplements (electrolytes, trace elements and vitamins)

See section 4.2. For specific admixtures, chemical and physical in-use stability has been demonstrated for 7 days (between 2°C and 8°C) followed by 48 hours at temperature not exceeding 25°C.

From a microbiological point of view, any admixture should be used immediately. If not used immediately, in-use storage times and conditions, after mixing and prior to use, are the responsibility of the user and would normally not be longer than 24 hours at 2°C to 8°C.

6.5 Nature and contents of container

The three-compartment bag is a multi-layer plastic bag. The inner (contact) layer of the bag is made of a blend of polyolefinic copolymers and is compatible with amino acid solutions, glucose solutions and lipid emulsions. Other layers are made of poly-ethylene vinyl acetate (EVA) and of copolyester.

The glucose compartment is fitted with an injection site to be used for addition of supplements. The amino acid compartment is fitted with an administration site for insertion of the spike of the infusion set.

The bag is packaged in an oxygen barrier overpouch which contains an oxygen absorber/oxygen indicator sachet.

Pack sizes

Formulation	Bag size				
PeriOlimel N4-600E	–	1000mL	1500mL	2000mL	2500mL
Olimel N5-860E	–	–	1500mL	2000mL	2500mL
Olimel N7-960	–	1000mL	1500mL	2000mL	–
Olimel N7-960E	–	1000mL	1500mL	2000mL	–
Olimel N9-840	–	1000mL	1500mL	2000mL	–
Olimel N9-840E	–	1000mL	1500mL	2000mL	–
Olimel N12-640	650mL	1000mL	1500mL	2000mL	–
Olimel N12-640E	650mL	1000mL	1500mL	2000mL	–

Note: Not all formulations and/or bag sizes may be marketed.

6.6 Special precautions for disposal

Any unused product or waste material should be disposed of in accordance with local requirements.

7 MEDICINE SCHEDULE

General Sale Medicine.

NEW ZEALAND DATA SHEET

8 SPONSOR

Olimel/PeriOlimel are distributed in New Zealand by:

Baxter Healthcare Ltd
33 Vestey Drive
Mt Wellington
Auckland 1060.

Baxter Healthcare Ltd
PO Box 14 062
Panmure
Auckland 1741

Phone (09) 574 2400.

Olimel/PeriOlimel are distributed in Australia by:

Baxter Healthcare Pty Ltd
1 Baxter Drive
Old Toongabbie, NSW 2146.

9 DATE OF FIRST APPROVAL

Date of publication in the New Zealand Gazette of consent to distribute the medicine:

PeriOlimel N4-600E, **Olimel** N5-860E

Olimel N7-960, **Olimel** N7-960E

Olimel N9-840, **Olimel** N9-840E

28 February 2013.

Olimel N12-640 and **Olimel** N12-640E

17 September 2020.

10 DATE OF REVISION OF THE TEXT

17 September 2020

SUMMARY TABLE OF CHANGES

Section changed	Summary of new information
ALL	Data Sheet updated to include information on Olimel N12-640 and Olimel N12-640E.

Based on Australian PI most recent amendment 23 October 2019 and CCDS402 2019JUL02.

Please refer to the Medsafe website (www.medsafe.govt.nz) for most recent data sheet.

Baxter, PeriOlimel, Olimel and Oliclinomel are trademarks of Baxter International Inc.

NEW ZEALAND DATA SHEET

APPENDIX 1

Composition per Litre of Reconstituted Emulsion								
Active substances	PeriOlimel N4-600E	Olimel N5-860E	Olimel N7-960E	Olimel N7-960	Olimel N9-840E	Olimel N9-840	Olimel N12- 640E	Olimel N12- 640
Refined olive oil + refined soybean oil*	30.00g	40.00g	40.00g	40.00g	40.00g	40.00g	35.00g	35.00g
L-Alanine	3.66g	4.76g	6.41g	6.41g	8.24g	8.24g	10.99g	10.99g
L-Arginine	2.48g	3.22g	4.34g	4.34g	5.58g	5.58g	7.44g	7.44g
L-Aspartic acid	0.73g	0.95g	1.28g	1.28g	1.65g	1.65g	2.20g	2.20g
L-Glutamic acid	1.26g	1.64g	2.21g	2.21g	2.84g	2.84g	3.79g	3.79g
Glycine	1.76g	2.28g	3.07g	3.07g	3.95g	3.95g	5.26g	5.26g
L-Histidine	1.51g	1.96g	2.64g	2.64g	3.40g	3.40g	4.53g	4.53g
L-Isoleucine	1.26g	1.64g	2.21g	2.21g	2.84g	2.84g	3.79g	3.79g
L-Leucine	1.76g	2.28g	3.07g	3.07g	3.95g	3.95g	5.26g	5.26g
L-Lysine acetate (equiv Lysine)	2.81g (1.99g)	3.65g (2.59g)	4.88g (3.48g)	4.88g (3.48g)	6.32g (4.48g)	6.32g (4.48g)	8.43g (5.97g)	8.43g (5.97g)
L-Methionine	1.26g	1.64g	2.21g	2.21g	2.84g	2.84g	3.79g	3.79g
L-Phenylalanine	1.76g	2.28g	3.07g	3.07g	3.95g	3.95g	5.26g	5.26g
L-Proline	1.51g	1.96g	2.64g	2.64g	3.40g	3.40g	4.53g	4.53g
L-Serine	1.00g	1.30g	1.75g	1.75g	2.25g	2.25g	3.00g	3.00g
L-Threonine	1.26g	1.64g	2.21g	2.21g	2.84g	2.84g	3.79g	3.79g
L-Tryptophan	0.42g	0.55g	0.74g	0.74g	0.95g	0.95g	1.26g	1.26g
L-Tyrosine	0.06g	0.08g	0.11g	0.11g	0.15g	0.15g	0.20g	0.20g
L-Valine	1.62g	2.10g	2.83g	2.83g	3.64g	3.64g	4.86g	4.86g
Sodium acetate, trihydrate	1.16g	1.50g	1.50g	--	1.50g	--	1.5g	--
Sodium glycerophosphate, hydrated	1.91g	3.67g	3.67g	--	3.67g	--	3.67g	--
Potassium chloride	1.19g	2.24g	2.24g	--	2.24g	--	2.24g	--
Magnesium chloride, hexahydrate	0.45g	0.81g	0.81g	--	0.81g	--	0.81g	--
Calcium chloride, dihydrate	0.30g	0.52g	0.52g	--	0.52g	--	0.52g	--
Glucose monohydrate (equiv to anhydrous glucose)	82.5g (75.0g)	126.5g (115.0g)	154.0g (140.0g)	154.0g (140.0g)	121.0g (110.0g)	121.0g (110.0g)	80.7g (73.3g)	80.7g (73.3g)

* Mixture of refined olive oil (approximately 80%) and refined soya-bean oil (approximately 20%), corresponding to a ratio essential fatty acids / total fatty acids of 20%.
The soya oil ingredient may contain ascorbyl palmitate as an antioxidant in the concentration $\leq 0.15\text{mg/g}$ of soya oil.

NEW ZEALAND DATA SHEET

After the contents of the three compartments have been mixed, the 3-in-1 admixture for each of the bag presentations provides the following:

For PeriOlimel N4-600E

	1000mL	1500mL	2000mL	2500mL
Nitrogen	4.0g	6.0g	8.0g	10.0g
Amino acids	25.3g	38.0g	50.6g	63.3g
Glucose monohydrate	82.5g	123.75g	165.0g	206.25g
Lipids	30.0g	45.0g	60.0g	75.0g
Energy:				
Total calories approx.	700kcal	1050kcal	1400kcal	1750kcal
Non-protein calories approx.	600kcal	900kcal	1200kcal	1500kcal
Glucose calories	300kcal	450kcal	600kcal	750kcal
Lipid calories⁽¹⁾	300kcal	450kcal	600kcal	750kcal
Non-protein calories / nitrogen ratio	150kcal/g	150kcal/g	150kcal/g	150kcal/g
Glucose / lipid calories ratio	50 / 50	50 / 50	50 / 50	50 / 50
Lipid / total calories	43%	43%	43%	43%
Electrolytes:				
Sodium	21.0mmol	31.5mmol	42.0mmol	52.5mmol
Potassium	16.0mmol	24.0mmol	32.0mmol	40.0mmol
Magnesium	2.2mmol	3.3mmol	4.4mmol	5.5mmol
Calcium	2.0mmol	3.0mmol	4.0mmol	5.0mmol
Phosphate⁽²⁾	8.5mmol	12.7mmol	17.0mmol	21.2mmol
Acetate	27.0mmol	41.0mmol	55.0mmol	69.0mmol
Chloride	24.0mmol	37.0mmol	49.0mmol	61.0mmol
pH	6.4	6.4	6.4	6.4
Osmolarity	760mOsm/L	760mOsm/L	760mOsm/L	760mOsm/L
(1) Include calories from egg lecithin (purified egg phosphatide)				
(2) Includes phosphate from lipid emulsion (egg phospholipids)				

NEW ZEALAND DATA SHEET

For Olimel N5-860E

	1500mL	2000mL	2500mL
Nitrogen	7.8g	10.4g	13.0g
Amino acids	49.4g	65.8g	82.3g
Glucose monohydrate	189.75g	253.0g	316.25g
Lipids	60.0g	80.0g	100.0g
Energy:			
Total calories	1490kcal	1980kcal	2480kcal
Non-protein calories	1290kcal	1720kcal	2150kcal
Glucose calories	690kcal	920kcal	1150kcal
Lipid calories (approx)⁽¹⁾	600kcal	800kcal	1000kcal
Non-protein calories / nitrogen ratio	165kcal/g	165kcal/g	165kcal/g
Glucose / lipid calories ratio	53 / 47	53 / 47	53 / 47
Lipid / total calories	47%	47%	47%
Electrolytes:			
Sodium	52.5mmol	70.0mmol	87.5mmol
Potassium	45.0mmol	60.0mmol	75.0mmol
Magnesium	6.0mmol	8.0mmol	10.0mmol
Calcium	5.3mmol	7.0mmol	8.8mmol
Phosphate⁽²⁾	22.5mmol	30.0mmol	37.5mmol
Acetate	55.0mmol	73.0mmol	91.0mmol
Chloride	68.0mmol	90.0mmol	113.0mmol
pH	6.4	6.4	6.4
Osmolarity	1120mOsm/L	1120mOsm/L	1120mOsm/L
(1) Include calories from egg lecithin (purified egg phosphatide)			
(2) Includes phosphate from lipid emulsion (egg phospholipids)			

NEW ZEALAND DATA SHEET

For Olimel N7-960

	1000mL	1500mL	2000mL
Nitrogen	7.0g	10.5g	14.0g
Amino acids	44.3g	66.4g	88.6g
Glucose monohydrate	154.0g	231.0g	308.0g
Lipids	40.0g	60.0g	80.0g
Energy:			
Total calories	1140kcal	1710kcal	2270kcal
Non-protein calories	960kcal	1440kcal	1920kcal
Glucose calories	560kcal	840kcal	1120kcal
Lipid calories⁽¹⁾	400kcal	600kcal	800kcal
Non-protein calories / nitrogen ratio	137kcal/g	137kcal/g	137kcal/g
Glucose / lipid calories ratio	58 / 42	58 / 42	58 / 42
Lipid / total calories	35%	35%	35%
Electrolytes:			
Phosphate⁽²⁾	3.0mmol	4.5mmol	6.0mmol
Acetate	31.0mmol	46.0mmol	62.0mmol
pH	6.4	6.4	6.4
Osmolarity	1220mOsm/L	1220mOsm/L	1220mOsm/L
(1) Include calories from egg lecithin (purified egg phosphatide)			
(2) Includes phosphate from lipid emulsion (egg phospholipids)			

For Olimel N7-960E

	1000mL	1500mL	2000mL
Nitrogen	7.0g	10.5g	14.0g
Amino acids	44.3g	66.4g	88.6g
Glucose monohydrate	154.0g	231.0g	308.0g
Lipids	40.0g	60.0g	80.0g
Energy:			
Total calories	1140kcal	1710kcal	2270kcal
Non-protein calories	960kcal	1440kcal	1920kcal
Glucose calories	560kcal	840kcal	1120kcal
Lipid calories⁽¹⁾	400kcal	600kcal	800kcal
Non-protein calories / nitrogen ratio	137kcal/g	137kcal/g	137kcal/g
Glucose / lipid calories ratio	58 / 42	58 / 42	58 / 42
Lipid / total calories	35%	35%	35%
Electrolytes:			
Sodium	35.0mmol	52.5mmol	70.0mmol
Potassium	30.0mmol	45.0mmol	60.0mmol
Magnesium	4.0mmol	6.0mmol	8.0mmol
Calcium	3.5mmol	5.3mmol	7.0mmol
Phosphate⁽²⁾	15.0mmol	22.5mmol	30.0mmol
Acetate	45.0mmol	67.0mmol	89.0mmol
Chloride	45.0mmol	68.0mmol	90.0mmol
pH	6.4	6.4	6.4
Osmolarity	1360mOsm/L	1360mOsm/L	1360mOsm/L
(1) Include calories from egg lecithin (purified egg phosphatide)			
(2) Includes phosphate from lipid emulsion (egg phospholipids)			

NEW ZEALAND DATA SHEET

For Olimel N9-840

	1000mL	1500mL	2000mL
Nitrogen	9.0g	13.5g	18.0g
Amino acids	56.9g	85.4g	113.9g
Glucose monohydrate	121.0g	181.5g	242.0g
Lipids	40.0g	60.0g	80.0g
Energy:			
Total calories	1070kcal	1600kcal	2140kcal
Non-protein calories	840kcal	1260kcal	1680kcal
Glucose calories	440kcal	660kcal	880kcal
Lipid calories⁽¹⁾	400kcal	600kcal	800kcal
Non-protein calories / nitrogen ratio	93kcal/g	93kcal/g	93kcal/g
Glucose / lipid calories ratio	52 / 48	52 / 48	52 / 48
Lipid / total calories	37%	37%	37%
Electrolytes:			
Phosphate⁽²⁾	3.0mmol	4.5mmol	6.0mmol
Acetate	40.0mmol	60.0mmol	80.0mmol
pH	6.4	6.4	6.4
Osmolarity	1170mOsm/L	1170mOsm/L	1170mOsm/L
(1) Include calories from egg lecithin (purified egg phosphatide)			
(2) Includes phosphate from lipid emulsion (egg phospholipids)			

For Olimel N9-840E

	1000mL	1500mL	2000mL
Nitrogen	9.0g	13.5g	18.0g
Amino acids	56.9g	85.4g	113.9g
Glucose monohydrate	121.0g	181.5g	242.0g
Lipids	40.0g	60.0g	80.0g
Energy:			
Total calories	1070kcal	1600kcal	2140kcal
Non-protein calories	840kcal	1260kcal	1680kcal
Glucose calories	440kcal	660kcal	880kcal
Lipid calories⁽¹⁾	400kcal	600kcal	800kcal
Non-protein calories / nitrogen ratio	93kcal/g	93kcal/g	93kcal/g
Glucose / lipid calories ratio	52 / 48	52 / 48	52 / 48
Lipid / total calories	37%	37%	37%
Electrolytes:			
Sodium	35.0mmol	52.5mmol	70.0mmol
Potassium	30.0mmol	45.0mmol	60.0mmol
Magnesium	4.0mmol	6.0mmol	8.0mmol
Calcium	3.5mmol	5.3mmol	7.0mmol
Phosphate⁽²⁾	15.0mmol	22.5mmol	30.0mmol
Acetate	54.0mmol	80.0mmol	107.0mmol
Chloride	45.0mmol	68.0mmol	90.0mmol
pH	6.4	6.4	6.4
Osmolarity	1310mOsm/L	1310mOsm/L	1310mOsm/L
(1) Include calories from egg lecithin(purified egg phosphatide)			
(2) Includes phosphate from lipid emulsion (egg phospholipids)			

NEW ZEALAND DATA SHEET

For Olimel N12-640

	1000mL	1500mL	2000mL	2500mL
Nitrogen	7.8g	12.0g	18.0g	24.0g
Amino acids	49.4g	75.9g	113.9g	151.9g
Glucose monohydrate	52.4g	80.7g	121.0g	161.3g
Lipids	22.8g	35.0g	52.5g	70.0g
Energy:				
Total calories approx.	620kcal	950kcal	1420kcal	1900kcal
Non-protein calories approx.	420kcal	640kcal	960kcal	1280kcal
Glucose calories	190kcal	290kcal	430kcal	580kcal
Lipid calories⁽¹⁾	230kcal	350kcal	520kcal	700kcal
Non-protein calories / nitrogen ratio	53kcal/g	53kcal/g	53kcal/g	53kcal/g
Glucose / lipid calories ratio	45 / 55	45 / 55	45 / 55	45 / 55
Lipid / total calories	37%	37%	37%	37%
Electrolytes:				
Phosphate⁽²⁾	1.7mmol	2.6mmol	3.9mmol	5.2mmol
Acetate	35mmol	54mmol	80mmol	107mmol
pH	6.4	6.4	6.4	6.4
Osmolarity	1130mOsm/L	1130mOsm/L	1130mOsm/L	1130mOsm/L
(1) Include calories from egg lecithin (purified egg phosphatide)				
(2) Includes phosphate from lipid emulsion (egg phospholipids)				

For Olimel N12-640E

	1000mL	1500mL	2000mL	2500mL
Nitrogen	7.8g	12.0g	18.0g	24.0g
Amino acids	49.4g	75.9g	113.9g	151.9g
Glucose monohydrate	22.8g	35.0g	52.5g	70.0g
Lipids	52.4g	80.7g	121.0g	161.3g
Energy:				
Total calories approx.	620kcal	950kcal	1420kcal	1900kcal
Non-protein calories approx.	420kcal	640kcal	960kcal	1280kcal
Glucose calories	190kcal	290kcal	430kcal	580kcal
Lipid calories⁽¹⁾	230kcal	350kcal	520kcal	700kcal
Non-protein calories/nitrogen ratio	53kcal/g	53kcal/g	53kcal/g	53kcal/g
Glucose / lipid calories ratio	45 / 55	45 / 55	45 / 55	45 / 55
Lipid / total calories	37%	37%	37%	37%
Electrolytes:				
Sodium	22.8mmol	35.0mmol	52.5mmol	70.0mmol
Potassium	19.5mmol	30.0mmol	45.0mmol	60.0mmol
Magnesium	2.6mmol	4.0mmol	6.0mmol	8.0mmol
Calcium	2.3mmol	3.5mmol	5.3mmol	7.0mmol
Phosphate⁽²⁾	9.5mmol	15.0mmol	21.9mmol	29.2mmol
Acetate	46mmol	70mmol	105mmol	140mmol
Chloride	30mmol	45mmol	68mmol	90mmol
pH	6.4	6.4	6.4	6.4
Osmolarity	1270mOsm/L	1270mOsm/L	1270mOsm/L	1270mOsm/L
(1) Include calories from egg lecithin (purified egg phosphatide)				
(2) Includes phosphate from lipid emulsion (egg phospholipids)				